arXiv:1207.1184v1 [cond-mat.stat-mech] 5 Jul 2012

Exact solution of a Levy walk model for anomalous heat transport

Abhishek Dhar,[1] Keiji Saito,[2] and Bernard Derrida[3]
[1] *International centre for theoretical sciences, TIFR, Bangalore 560012, India*
[2] *Department of Physics, Keio University, Yokohama 223-8522, Japan*
[3] *Laboratoire de Physique Statistique, Ecole Normale Supérieure, UPMC,*
Université Paris Diderot, CNRS, 24 rue Lhomond, 75231 Paris cedex 05, France
(Dated: July 6, 2012)

The Levy-walk model is known to provide a good description of anomalous heat conduction in one-dimensional systems. In this model the heat carriers execute Levy-walks instead of normal diffusion as expected in systems where Fourier's law holds. Here we calculate exactly the average heat current, the large deviation function of its fluctuations and the temperature profile of the Levy-walk model maintained in a steady state by contact with two heat baths (the open geometry). We find that the current is nonlocally connected to the temperature gradient. As observed in recent simulations of mechanical models, all the cumulants of the current fluctuations have the same system-size dependence in the open geometry. The case of the ring geometry is also discussed.

PACS numbers:

Introduction.— Fourier's law of heat conduction (in one dimension) states that

$$J(x,t) = -\kappa \frac{dT(x,t)}{dx}, \tag{1}$$

where $T(x,t)$, $J(x,t)$ are the local temperature and heat current density fields and κ is the thermal conductivity. Based on results obtained from a large number of numerical simulations and various analytical approaches it is now believed that Fourier's law is not valid in one and two dimensional mechanical systems (in particular models where momentum is conserved) and heat conduction is anomalous [1–3]. Anomalous behavior in heat conduction is not only a theoretical issue but recently of experimental relevance in several low-dimensional materials [4, 5]. Indicators of the anomaly include — (i) in steady states the dependence of the heat current J on system size L shows the scaling behaviour $J \sim L^{\alpha-1}$ with $\alpha > 0$, (ii) the temperature profiles across systems in nonequilibrium steady states are found to be nonlinear, even for very small applied temperature differences and, (iii) the spreading of heat pulses in anharmonic chains is super-diffusive.

The microscopic basis of Fourier's law lies in the fact that the carriers of heat in a system execute random walks. The simplest "derivation" of Fourier's law, from kinetic theory, is based on this picture and leads to an expression of the conductivity of a material in terms of the mean free path ℓ, mean velocity v and specific heat capacity c of the heat carriers: $\kappa = cv\ell$ (in one dimensions). The breakdown of Fourier's law thus also implies a breakdown of the random walk picture and of the diffusion equation

$$\frac{\partial T(x,t)}{\partial t} = \frac{\kappa}{c}\frac{\partial^2 T(x,t)}{\partial x^2}~, \tag{2}$$

which describes time-dependent heat transfer (assuming κ has no temperature-dependence). A number of recent studies indicate that a good description of anomalous heat conduction in one dimensional systems is obtained by modeling the motion of the heat carriers as Levy random walks instead of simple random walks [6–9]. Numerical studies show that the spreading of heat pulses and the form of steady state temperature profiles can be correctly modeled by means of the Levy walk. In another study of a model with stochastic dynamics, it has been shown, starting from a Boltzmann-equation approach, that the temperature satisfies a fractional diffusion equation corresponding to a Levy stable process [10].

While some analytical understanding has been achieved [6] it is desirable to further develop the Levy walk theory for anomalous heat conductivity so that (i) one can use it in the way as one uses Eqs. (1,2) for normal diffusion and (ii) one has a clearer idea of the range of applicability of the model. In this Letter we present several exact results for steady state heat transport in the Levy walk model in one dimension. For setting up the steady state we follow the idea in [9] of connecting two infinite reservoirs at different temperatures to the system and consider a version where space and time are taken to be continuous.

Our exact results provide several interesting physical perspective on anomalous heat transport. The analytic solution of the particle-density profile exhibits the nonlinear (and singular) form typical of temperature profiles in $1D$ systems [2, 3]. The steady state current has the power law dependence on the system-size, characteristic of anomalous diffusion. In contrast to (1) for normal heat transport, it is non-locally connected to the temperature gradient. We derive the exact cumulant generating function of current for the open geometry. Our results show that all cumulants of the integrated current have the same system-size dependence as the average current. This is consistent with recent numerical results of heat conduction in hard particle systems [11] and therefore strongly suggests that the Levy-walk model gives a good description of anomalous heat conduction not only for the average current but also for current fluctuations. We also show that the cumulant generating function is consistent with the additivity principle expression [19, 20]. For the ring geometry, also, the size dependence of the cumulants obtained in simulations [11] can be recovered by introducing a size dependent cut-off in the distribution of times of the Levy-walk model.

Levy diffusion on the infinite line.— In the simplest description we think of energy in the system as being effectively mediated by particles performing Levy walks. Each particle carries a single quantum of energy. Therefore the local energy density and energy current at any point are directly proportional to the particle density and current respectively. The precise definition of the Levy walk model that we consider here is as follows. For a single particle each step of the walk consists in choosing a time of flight τ from a given distribution $\phi(\tau)$ and then moving it at speed v over a distance $x = v\tau$ in either direction, with equal probability. Let us define $P(x,t)dx$ as the probability that the particle is in the interval $(x, x+dx)$ at time t. Thus $P(x,t)$ includes events where the particle is crossing the interval $(x, x+dx)$. If a particle starts at the origin at time $t = 0$, the probability $P(x,t)$ satisfies

$$\begin{aligned} P(x,t) &= \frac{1}{2}\psi(t)\delta(|x|-vt) \qquad (3)\\ &+\frac{1}{2}\int_0^t d\tau\phi(\tau)[P(x-v\tau,t-\tau)+P(x+v\tau,t-\tau)] \ , \end{aligned}$$

where $\psi(\tau) = \int_\tau^\infty d\tau' \ \phi(\tau')$ is the probability of choosing a time of flight $\geq \tau$. The Fourier-Laplace transform of $P(x,t)$ can be calculated from (3) in terms of the Laplace transform of the distribution of flight times $\phi(\tau)$ and this gives the time dependence of all the cumulants $\langle x^n\rangle_c$ of the position x at time t.

If the first moments of the flight times $\langle\tau^n\rangle$ were finite, the motion would be diffusive. One would then get from (3) for the first cumulants of x at large t

$$\frac{\langle x^2\rangle_c}{v^2 \ t} \simeq \frac{\langle\tau^2\rangle}{\langle\tau\rangle} \ ; \ \ \frac{\langle x^4\rangle_c}{v^4 \ t} \simeq 3\frac{\langle\tau^2\rangle^3}{\langle\tau\rangle^3} - 6\frac{\langle\tau^2\rangle\langle\tau^3\rangle}{\langle\tau\rangle^2} + \frac{\langle\tau^4\rangle}{\langle\tau\rangle} \ . \quad (4)$$

Here we will consider time-of-flight distributions with a power law decay at large time

$$\phi(\tau) \simeq A \ \tau^{-\beta-1} \ , \ \ 1 < \beta < 2 \ , \qquad (5)$$

(for example $\phi(\tau) = \beta/t_o/(1+\tau/t_o)^{\beta+1}$ which is used for the data shown in figure 1). For this range of β the mean flight time $\langle\tau\rangle = \int_0^\infty d\tau \ \tau \ \phi(\tau)$ is finite but $\langle\tau^2\rangle = \infty$. Using again the Fourier-Laplace transform of (3) one gets for large t

$$\langle x^2\rangle_c \simeq \frac{2 \ A \ v^2}{(3-\beta)(2-\beta)\beta \ \langle\tau\rangle} t^\gamma, \qquad \gamma = 3-\beta \ , \quad (6)$$

$$\langle x^4\rangle_c \simeq \frac{4 \ A \ v^4}{(5-\beta)(4-\beta)\beta \ \langle\tau\rangle} t^{\gamma+2} \ . \quad (7)$$

We see that for $1 < \beta < 2$ the motion is superdiffusive [12, 13]. The solution of (3) corresponds to a pulse whose central region is a Levy-stable distribution with a scaling $x \sim t^{1/\beta}$, has ballistic peaks of magnitude $t^{1-\beta}$ at $x = \pm vt$ and vanishes outside this [8].

From the evolution equation (3), one can also see that a density fluctuation of wave number k relaxes exponentially with a time constant t^* obtained from the solution of $\int_0^\infty d\tau\phi(\tau)\cos(kv\tau)\exp[\tau/t^*] = 1$. For $\phi(\tau)$ of the form (5) one gets for small k

$$1/t^* \simeq A\Gamma(-\beta)\cos(\pi(1-\beta/2)) \ (kv)^\beta/\langle\tau\rangle \qquad (8)$$

whereas when $\phi(\tau)$ has a finite $\langle\tau^2\rangle$, the regime is diffusive with $t^* \sim k^2$.

Levy diffusion in a finite system connected to infinite reservoirs.— Let us now turn to the more interesting case of the open geometry where the system is a finite segment between $(0, L)$ connected on its two sides to reservoirs. In addition to the probability density $P(x,t)$ let us also define the quantity $Q(x,t)dxdt$ as the probability that a particle has precisely landed in the interval $(x, x+dx)$ during the time interval $(t, t+dt)$. Note that at any given time, a particle could either have landed at a point x or could be passing over the point. We need to set up the correct boundary conditions required to construct a nonequilibrium current carrying steady state. To do so we identify the region $x \leq 0$ with the left reservoir and the region $x \geq L$ with the right reservoir. We set $Q(x,t) = Q_l$ for points on the left reservoir and $Q(x,t) = Q_r$ for those on the right. In the steady state we have $Q(x,t) = Q(x)$ and $P(x,t) = P(x)$, and they satisfy

$$\begin{aligned} Q(x) - \int_0^L dx' \ \frac{1}{2v}\phi(|x-x'|/v) \ Q(x') & \\ = \frac{Q_l}{2}\psi(x/v) + \frac{Q_r}{2}\psi[(L-x)/v] \ , & \qquad (9)\\ P(x) = \int_0^L dx' \frac{1}{2v}\psi(|x-x'|/v) \ Q(x') & \\ + \frac{Q_l}{2}\chi(x/v) + \frac{Q_r}{2}\chi[(L-x)/v] \ . & \qquad (10) \end{aligned}$$

where, as in (3), $\psi(t) = \int_t^\infty d\tau \ \ \phi(\tau)$ and $\chi(t) = \int_t^\infty d\tau\psi(\tau)$. The problem is closely related to the escape probability [14] of a Levy walker on the interval $(0, L)$: if $H(x)$ is the probability that a Levy walker starting at position x will first hit the left reservoir (i.e. the region $x < 0$) before it hits the right reservoir (i.e. $x > L$) it is



FIG. 1: Plot of $Q(x)$ for different system sizes, for the Levy walk with $\beta = 1.5, Q_l = 1.0, Q_r = 0.5$. The data are obtained by solving Eqs. (9,10) with discretized space. The distribution of flight times is $\phi(\tau) = \beta/t_o/(1+\tau/t_o)^{\beta+1}$. The inset shows $P(x)/Q(x)$, which converges to $\chi(0)$ in the limit of $L \to \infty$.

easy to see that $H(x)$ satisfies

$$H(x) = \int_0^L dx' \ \frac{1}{2v}\phi(|x-x'|/v) \ H(x') + \frac{1}{2}\psi(x/v) \ , \quad (11)$$

and from (9), one can see that $Q(x) = (Q_l - Q_r)H(x) + Q_r$.

By following the same mathematical steps as in [14] to study equations such as (9) or (11), one can show that, in the large L limit, the solution $Q(x)$ of (9) (and $H(x)$ of (11)) satisfy

$$\int_0^L dx' \psi(|x-x'|/v) \ \mathrm{Sgn}(x-x') Q'(x') = 0 \ . \quad (12)$$

with $Q(0) = Q_l$ and $Q(L) = Q_r$ (and $H(0) = 1$ and $H(1) = 0$ for (11)) with a solution of (12), for a $\phi(\tau)$ decaying as in (5), which satisfies

$$Q'(x) = -B[x(L-x)]^{\beta/2-1} \ . \quad (13)$$

Integrating this and imposing the boundary conditions $Q(0) = Q_l$ and $Q(L) = Q_r$, one obtains the constant $B = (Q_l - Q_r) \ \Gamma(\beta)/\Gamma(\beta/2)^2 \ L^{1-\beta}$ and therefore $Q(x)$. One can also notice that in (10) the r.h.s. is dominated, for large L, by the range $|x'-x| \ll L$ and therefore

$$P(x) = \chi(0)Q(x) = \langle\tau\rangle Q(x) \ . \quad (14)$$

In Fig. (1), we compare numerical results obtained by solving Eqs. (9,10) with the exact results of Eqs. (13,14). The profiles are nonlinear and look similar to those observed for temperature profiles in 1D heat conduction [2, 3].

We next discuss the current. The steady state current $J(x)$ at position x is given by

$$J(x) = \frac{1}{2}\int_{-\infty}^{\infty} dx' \ Q(x-x') \ \mathrm{Sgn}(x') \ \psi(|x'|/v) \ , \quad (15)$$

which can be interpreted as the difference between the flow from left to right and from right to left. After a partial integration and using the fact that $Q(0) = Q_l$ and $Q(L) = Q_r$, one gets

$$J(x) \ = \ -\frac{v}{2}\int_0^L dx' \chi(|x-x'|/v) \ Q'(x') \ . \quad (16)$$

We note that $dJ/dx = 0$ gives Eq. (12) and so the current is independent of x, as expected. Evaluating the current at $x = 0$ and using Eq. (13), we get for large L

$$J \simeq (Q_l - Q_r) \ \frac{A \ v^\beta \ \Gamma(\beta) \ \Gamma(1-\frac{\beta}{2})}{2 \ \beta(\beta-1) \ \Gamma(\frac{\beta}{2})} \ L^{\alpha-1}, \quad \alpha = 2-\beta. \quad (17)$$

From Eq. (6) we then get the relation $\alpha = \gamma - 1$, between the conductivity exponent of anomalous transport and the exponent for Levy-walk diffusion. This relation for Levy diffusion was noted in [6], numerically observed in $1D$ heat conduction models [8, 15] and a derivation based on linear response theory has recently been proposed [16].

In the large L limit by using Eq. (14) in Eq. (16) we obtain

$$J \ = \ -\frac{v}{2\langle\tau\rangle}\int_0^L dx' \chi(|x-x'|/v) P'(x') \ . \quad (18)$$

This is the analogue of Fourier's Law Eq. (1) in the case of normal heat conduction and shows that the current is nonlocally connected to the temperature gradient.

Current fluctuations in the open system.— In the rest of this letter, we discuss current fluctuations.

The only contribution to the current through the systems is from walkers injected (at rate p_L) from the left reservoir which end up (either after a non stop flight or a non direct flight) into the right reservoir or walkers injected (at rate p_R) from the right reservoir which end up into the left reservoir. In the case $Q_l = 1$ and $Q_r = 0$ let $\mathcal{P}$ be the rate at which walkers, which will end up into the right reservoir, are injected from the left reservoir. For general Q_l and Q_r, because the walkers are independent and because they have no preferred direction, one has $p_L = Q_l\mathcal{P}$ and $p_R = Q_r\mathcal{P}$. We can then write the characteristic function of the integrated current $\mathcal{Q} = \int_0^\tau dt J(t)$ in time τ,

$$\begin{aligned} Z(\lambda) &= \langle e^{\lambda\mathcal{Q}}\rangle = \Pi_t\langle e^{\lambda J(t)dt}\rangle \\ &= \Pi_t[(1 - p_L dt - p_R dt) + e^\lambda p_L dt + e^{-\lambda} p_R dt] \\ &= e^{\mu(\lambda)\tau} \ , \\ \text{where} \ \ \mu(\lambda) &= p_L(e^\lambda - 1) + p_R(e^{-\lambda} - 1) \ . \end{aligned} \quad (19)$$

Hence we get for the cumulant generating function of $\mathcal{Q}$

$$\mu(\lambda) = J \ \frac{Q_l(e^\lambda - 1) + Q_r(e^{-\lambda} - 1)}{Q_l - Q_r} \ , \quad (20)$$

where $J = (Q_l - Q_r)\mathcal{P}$. We can check that $\mu'(\lambda = 0)$ gives the current J. Since in Eq.(20) the system-size occurs only through the average current J, all cumulants of $\mathcal{Q}$ have the same size dependence as the average current. This is consistent with recent numerical simulations of the 1D hard-point alternate mass gas [11]. Not surprisingly we also note that the following fluctuation theorem symmetry relation [17] is satisfied:

$$\mu(\lambda) = \mu[-\lambda - (\ln Q_l - \ln Q_r)] \ . \tag{21}$$

Current fluctuations in ring geometry.— In the ring geometry, the system consists of a fixed number N of particles which perform independent Levy walks on a ring of length L. In the steady state, the density of particles is uniform. As the walkers are independent, the cumulants of the integrated current $\mathcal{Q}$ are related to those of the displacement $x(t)$ of a single walker on the infinite line (in the steady state)

$$\langle \mathcal{Q}^n \rangle_c \sim \frac{N}{L^n} \langle x(t)^n \rangle_c = \frac{\rho}{L^{n-1}} \langle x(t)^n \rangle_c \tag{22}$$

where ρ is the density on the ring. If the walkers perform on the ring the same Levy walks as on the infinite line, the cumulants of $x(t)$ and therefore those of $\mathcal{Q}$ grow, as in (6,7), faster than linearly with time (same exponent but a different prefactor as, on the ring, the walker is in its steady state rather than starting a flight at $t = 0$).

On the other hand if one introduces a cut-off time $\tau_L \sim L^\delta$ in the distribution $\phi(\tau)$ (for example by arguing that τ_L should be of the order of t^*, the relaxation time corresponding to the shortest wave number on the ring $k = 2\pi/L$, implying from (8) that $\delta = \beta$; one could alternatively argue that, as for the open geometry all the length of the flights cannnot exceed the system size and therefore $\delta = 1$). With such a cut-off τ_L, the cumulants of $\mathcal{Q}$ would grow linearly in time (4,23), with an amplitude which depends on the system size and on the cumulant considered

$$\frac{\langle \mathcal{Q}^2 \rangle_c}{t} \sim L^{(2-\beta)\delta - 1} \quad ; \quad \frac{\langle \mathcal{Q}^4 \rangle_c}{t} \sim L^{(4-\beta)\delta - 3} \tag{23}$$

This cut-off time τ_L gives therefore a possible explanation for the behavior seen in simulations on the ring of hard-point alternate gas in figure 3 of [11], where the cumulants grow linearly in time with different system size dependence ($\langle \mathcal{Q}^2 \rangle_c / t \sim L^{-0.5}$ and $\langle \mathcal{Q}^4 \rangle_c / t \sim L^{0.5}$). Then one gets from (23) $\beta \sim 5/3$ and $\delta \sim 3/2$ leading through (17) to a value $\alpha = 1/3$ for the anomalous Fourier's law of the hard-point alternate gas in the open geometry consistent with most of the simulations done so far [3, 11, 18] for this system . How to predict the values of β and δ from first principles in Hamiltoninan systems is of course an interesting open question.

Discussion.— In this work we have studied the Levy diffusion model of anamalous heat transport. We have computed the average current (17), the energy profile (13,14) and the large deviation function of the integrated current (19) , in the open geometry, i.e. when the system is connected at its two ends to reservoirs. One remarkable result is that all the cumulants of the integrated current have the same anomalous size dependence. We have also proposed a simple possibility to explain the size dependence of the cumulants for the ring geometry.

An interesting observation is that the generating function of the integrated current in Eq. (20) matches exactly with the formula obtained from the additivity principle (AP) [19] which gives an expression for $\mu(\lambda)$ in terms of the the conductivity D and equilibrium current fluctuations σ defined respectively as

$$\begin{aligned} D(Q) &= \lim_{\Delta Q \to 0} LJ/\Delta Q \ , \\ \sigma(Q) &= L \lim_{t \to \infty} \langle \mathcal{Q}^2 \rangle / t \ . \end{aligned} \tag{24}$$

The expression for $\mu(\lambda)$ from AP is

$$\begin{aligned} \mu_{AP}(\lambda) &= -\frac{K}{L} \left[\int_{Q_r}^{Q_l} dQ \frac{D(Q)}{\sqrt{1 + 2K\sigma(Q)}} \right]^2 , \\ \text{with } \lambda &= \int_{Q_r}^{Q_l} dQ \frac{D(Q)}{\sigma(Q)} \left[\frac{1}{\sqrt{1 + 2K\sigma(Q)}} - 1 \right] . \end{aligned} \tag{25}$$

(this is a parametric expression: as K varies, μ and λ vary). From our exact results for $\mu(\lambda)$ we find $D = Lp$ and $\sigma = L\ \mu''(\lambda = 0) = 2DQ$. Using these in Eq. (25) and after explicitly performing the integrals we find $\mu_{AP}(\lambda) = \mu(\lambda)$. This result is somewhat surprising since the additivity principle is expected normally to hold for diffusive systems (here D and σ have a L-dependence, whereas in usual diffusive systems they don't). This reminds us of the recent numerical study on the validity of the additivity principle in mass-disordered harmonic crystals where, even in the anomalous regime, the additivity principle expression is found to be accurate [20].

We thank NORDITA where this work was initiated. AD and KS thank S. Denisov for many useful discussions. AD thanks DST for support through the Swarnajayanti fellowship. KS was supported by MEXT (23740289).

[1] F. Bonetto, J.L. Lebowitz, and L. Rey-Bellet, in *Mathematical Physics 2000*, edited by A. Fokas et. al. (Imperial College Press, London, 2000), p. 128.
[2] S. Lepri, R. Livi, and A. Politi, Phys. Rep. **377**, 1 (2003).
[3] A. Dhar, Adv. Phys. **57**, 457 (2008).
[4] C. W. Chang *et al* , Phys. Rev. Lett. **101**, 075903 (2008).
[5] D.L. Nika *et al* , Appl. Phys. Lett. **94**, 203103 (2009).
[6] S. Denisov, J. Klafter and M. Urbakh, Phys. Rev. Lett. **91**, 194301 (2003).
[7] R. Metzler and I. M. Sokolov, Phys. Rev. Lett. **92**, 089401 (2004).

[8] P. Cipriani, S. Denisov, and A. Politi, Phys. Rev. Lett. **94**, 244301 (2005); V. Zaburdaev, S. Denisov, and P. Hänggi, Phys. Rev. Lett. **106**, 180601 (2011).

[9] S. Lepri and A. Politi, Phys. Rev. E **83**, 030107 (2011).

[10] M. Jara, T. Komorowski, and S. Olla, Ann. App. Prob. **19**, 2270 (2009).

[11] E. Brunet, B. Derrida, A. Gerschenfeld, Europhys. Lett. **90**, 20004 (2010).

[12] G. Zumofen and J. Klafter, Phys. Rev. E **47**, 851 (1993).

[13] R. Metzler and A. Compte, Physica A **268**, 454 (1999).

[14] S. V. Buldyrev, S. Havlin, A. Ya. Kazakov, M. G. E. da Luz , E. P. Raposo, H. E. Stanley, and G. M. Viswanathan, Phys. Rev. E **64**, 041108 (2001).

[15] H. Zhao, Phys. Rev. Lett. **96**, 140602 (2006).

[16] S. Liu, N. Li, J. Ren, and B. Li, arXiv:1103.2835.

[17] G. Gallavotti and E. G. D. Cohen, Phys. Rev. Lett. **74**, 2694 (1995); J.L Lebowitz and H. Spohn, J. Stat. Phys., **95**, 333 (1999).

[18] P. Grassberger, W. Nadler and L. Yang, Phys. Rev. Lett. **89**, 18061 (2002).

[19] T. Bodineau and B. Derrida, Phys. Rev. Lett. **92**, 180601 (2004); T. Bodineau and B. Derrida, C. R. Physique **8**, 540 (2007); L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio and C. Landim, Phys. Rev. Lett. **94**, 030601 (2005); P.I. Hurtado and P.L. Garrido, Phys. Rev. Lett. **102**, 250601 (2009).

[20] K. Saito and A. Dhar, Phys. Rev. Lett. **107**, 250601 (2011).